

## **Delirium Management: Evidence For Change And Future Directions**

E. Wesley Ely: Identification of Delirium in the ICU: Practical Points and Lessons Learned

Gregory Margolin: Protocols for Prevention and Management of Delirium in Different Patient and ICU Types

Richard R. Riker: Alpha-2 Agonists Versus GABA Agonists: Should We Change Standard Practice?

### References

1. American Psychiatric Association. *Diagnostic and Statistical Manual of Mental Disorders DSM-IV-TR Fourth Edition*. 4<sup>th</sup> ed. Arlington, VA: American Psychiatric Publishing, Inc; 2000.
2. Plaschke K, et al. Comparison of the confusion assessment method for the intensive care unit (CAM-ICU) with the Intensive Care Delirium Screening Checklist (ICDSC) for delirium in critical care patients gives high agreement rate(S). *Intensive Care Med*. 2008;34:431-436.
3. Delirium Assessment. ICU Delirium and Cognitive Impairment Study Group Web site. <http://www.icudelirium.org/delirium/training-pages/del.assessment.html> Accessed March 12, 2009.
4. Sommer BR, et al. Is dopamine administration possibly a risk factor for delirium? *Crit Care Med*. 2002;30:1508-1511.
5. Friese RS, Diaz-Arrastia R, et al. Quantity and quality of sleep in the surgical intensive care unit: are our patients sleeping? *J Trauma*. 2007;63(6):1210-1214.
6. Meagher DJ, et al. Relationship between symptoms and motoric subtype of delirium. *J Neuropsychiatry Clin Neurosci*. 2000;12:51-56.
7. Maldonado JR. Delirium in the acute care setting: characteristics, diagnosis and treatment. *Crit Care Clin*. 2008;24(4):657-722, vii. Review.

8. Pandharipande P, et al. Motoric subtypes of delirium in mechanically ventilated surgical and trauma intensive care unit patients. *Intensive Care Med.* 2007;33:1726-1731.
9. Jacobi J, Fraser GL, Coursin DB; Task Force of the American College of Critical Care Medicine (ACCM) of the Society of Critical Care Medicine (SCCM), American Society of Health-System Pharmacists (ASHP), American College of Chest Physicians. Clinical practice guidelines for the sustained use of sedatives and analgesics in the critically ill adult. *Crit Care Med.* 2002;30:119-141.
10. Gagnon B, et al. Methylphenidate hydrochloride improves cognitive function in patients with advanced cancer and hypoactive delirium: a prospective clinical study. *J Psychiatry Neurosci.* 2005;30:100-107.
11. Gold JA, Rimal B, Nolan A, Nelson LS. A strategy of escalating doses of benzodiazepines and Phenobarbital administration reduces the need for mechanical ventilation in deliria tremens. *Crit Care Med.* 2007;35:724-730.
12. Puntillo KA, et al. Patients' perceptions and responses to procedural pain: results from Thunder Project II. *Am J Crit Care.* 2001;10:238-251.
13. Pandharipande P, et al. Lorazepam is an independent risk factor for transitioning to delirium in intensive care unit patients. *Anesthesiology.* 2006;104:21-26.
14. Breen D, et al. Decreased duration of mechanical ventilation when comparing analgesia-based sedation using remifentanyl with standard hypnotic-based sedation for up to 10 days in intensive care unit patients: a randomized trial [ISRCTN47583497]. *Crit Care.* 2005;9:R200-210.
15. Pandharipande P, et al. Effect of sedation with dexmedetomidine vs lorazepam on acute brain dysfunction in mechanically ventilated patients: the MENDS randomized controlled trial. *JAMA.* 2007;298:2644-2653.
16. Riker RR, et al. Dexmedetomidine vs. midazolam for sedation of critically ill patients: a randomized trial. *JAMA.* 2009;301:489-499.

